arXiv:1106.4777v1 [math.GT] 23 Jun 2011

AUTOMORPHISMS OF THE 3-SPHERE THAT PRESERVE SPATIAL GRAPHS AND HANDLEBODY-KNOTS

YUYA KODA

Abstract. We consider the group of isotopy classes of automorphisms of S^3 that preserve a spatial graph or a handlebody-knot embedded in it. We prove that the group is finitely presented for an arbitrary spatial graph or a reducible handlebody-knot of genus two. We also prove that the groups for "most" irreducible handlebody-knots are finite.

2010 Mathematics Subject Classification: 57M15; 20F05, 57M25.
Keywords: spatial graph, handlebody, mapping class group.

Introduction

In the present paper, we consider two natural generalizations of the study of knots, the studies of spatial graphs and handlebody-knots.

By a *graph* G we shall mean a finite graph, possibly with loops and repeated edges. We allow the case where G has no vertices, that is, a component which is a loop. We consider a graph as a topological space in the usual way. A *spatial graph* Γ is a graph embedded in a 3-manifold M. Two spatial graphs are said to be equivalent if one can be transformed into the other by an ambient isotopy of the 3-manifold. It is clear that the theory of spatial knots is a generalization of the classical knot theory. Also, it is remarkable that a spatial trivalent graph appears as a singular locus of a 3-orbifold, see e.g. [8]. Moreover, most pairs of 3-manifolds and spatial trivalent graphs in them admit hyperbolic structures with parabolic meridians, that is, the complements of the spatial graphs carry metrics of constant sectional curvature -1 which complete to manifolds with non-compact geodesic boundary having toric cusps at the knot components of the graph, annular cusps at the meridians of the edges, and geodesic 3-punctured boundary spheres at the vertices, see e.g. [16]. Note that such structures are also useful when we study orbifold structures.

A *handlebody* is an orientable 3-manifold with boundary containing pairwise disjoint, properly embedded disks such that the manifold resulting from cutting along the disks is a 3-ball. A *handlebody-knot* is a handlebody embedded in a 3-manifold. Two handlebody-knots are said to be equivalent if one can be transformed into the other by an ambient isotopy of the 3-manifold. Handlebody-knot theory is another natural generalization of the classical knot theory, since a regular neighborhood of a knot determines a genus one handlebody-knot in a 3-manifold. A handlebody-knot also naturally appears in the theory of 3-manifolds, since every closed orientable 3-manifold admits a decomposition

---

Date:

into two handlebodies called a Heegaard splitting and such handlebodies can be regarded as simplest handlebody-knots in the 3-manifold. Moreover, such an object appears in the theory of hyperbolic structures with geodesic boundary, finiteness properties of invariants of 3-manifolds, etc.

A regular neighborhood of a spatial graph naturally be regarded as a handlebody-knot, however, regular neighborhoods of many different spatial graphs define the same handlebody-knot. For example, a regular neighborhood of the union of a tunnel number one knot and its unknotting tunnel is always the trivial handlebody-knot of genus two. Two spatial trivalent graphs define the same handlebody-knot if and only if one can be transformed into the other by a finite sequence of IH-moves (or Whitehead moves) for spatial graphs and ambient isotopies of the 3-manifold, see [18]. Here, an IH-move for a spatial graph is a replacement of a subspace shaped like the letter I in the spatial graph by a space shaped like the letter H.

For a subspace N of a compact orientable 3-manifold M, the mapping class group $\mathcal{MCG}(M, N)$ is defined to be the group of isotopy classes of the automorphisms of M that preserve N. We call the group $\mathcal{MCG}(M, N)$ the *symmetry group* of N. The aim of the present paper is to investigate whether the symmetry group $\mathcal{MCG}(M, N)$ is finitely presented when $M = S^3$ and N is a spatial graph or a handlebody-knot in S^3.

Recall the case where M is the 3-sphere S^3 and $N = K$ is a knot in S^3. Then the symmetry group $\mathcal{MCG}(S^3, K)$ of K has been widely studied by various authors, see [21]. In this case, the group $\mathcal{MCG}(S^3, N(K))$ is isomorphic to the mapping class group of $\overline{S^3 \setminus N(K)}$ due to a celebrated result of Gordon and Luecke [14], which states that if two knots in S^3 have homeomorphic complements then the homeomorphism between the two complements extends to an automorphism of S^3. It follows that the symmetry group $\mathcal{MCG}(S^3, N(K))$ is finitely presented due to the theory of mapping class groups of irreducible orientable sufficiently large 3-manifolds by McCullough [23]. Note, in contrast, that it is known that handlebody-knots of arbitrary genera are not determined by their complements, see [24, 25].

When V is a genus g unknotted handlebody in the 3-sphere S^3, the group $\mathcal{MCG}_+(S^3, V)$ is called the *Goeritz group* of genus g. In the case of genus two, it was shown by Goeritz [12] in 1933 that $\mathcal{MCG}_+(S^3, V)$ is finitely generated. Scharlemann [26] gave a modern, concise proof of this result, and Akbas [1] and Cho [5] independently gave a finite presentation of $\mathcal{MCG}_+(S^3, V)$. However, in the case of arbitrary genus, the existence of such generating sets still remains unknown. The best general reference here is the introduction of [5]. It is worth pointing out that finding a generating set of the symmetry groups of the unknotted handlebodies are very important for both the theory of handlebody-knots and that of Heegaard splittings.

The paper is organized as follows. In Section 1, we set up notation and terminology. Then we review some of the standard facts and histories on the symmetry groups. In Section 2, we discuss the case where M is the 3-sphere S^3 and $N = \Gamma$ is a spatial graph in S^3. Our first main result, Theorem 2.10, states that the symmetry group $\mathcal{MCG}(S^3, \Gamma)$ is finitely presented for an arbitrary spatial graph Γ in S^3. This is a generalization of a known result for knots to spatial graphs. From Section 3, we consider the symmetry groups of handlebody-knots. Note that we have $\mathcal{MCG}(S^3, K) \cong \mathcal{MCG}(S^3, N(K))$ for a knot $K \subset S^3$ by the uniqueness of a regular neighborhood. This implies that the theory of the symmetry groups of handlebody-knots is another generalization of that of knots. In Section 3, we consider the group $\mathcal{MCG}(S^3, V)$, where V is a reducible handlebody-knot of genus two. We define the complex $\mathcal{P}(V)$ of primitive disks, which is a subcomplex of the disk complex of a handlebody V. The construction is similar in spirit to the primitive disk complex for the genus two handlebody standardly embedded in S^3, see [5]. We prove that the complex $\mathcal{P}(V)$ is contractible and 1-dimensional, i.e. a tree, on which $\mathcal{MCG}(S^3, V)$ acts simplicially. We then conclude in Theorem 3.13 that the symmetry groups of reducible handlebody-knots of genus two are finitely presented due to the Bass-Serre theory of groups acting on graphs. It is worth noting that Cho [6] recently proved that the primitive disk complex of the unique genus two Heegaard splitting of a lens space is 1-dimensional or 2-dimensional, depending on the lens space. In Section 4, we provide a sufficient condition for the symmetry group of a irreducible handlebody-knots of genus two being finitely presented. It is highly expected that the symmetry groups of all irreducible handlebody-knots of genus two are finitely presented. We prove "most" of them are, in fact, finite groups, however, the result in the section is far from being conclusive.

Notation. Let X be a subset of a given topological space or a manifold Y. Throughout the paper, we will denote the interior of X by $\operatorname{Int} X$, the closure of X by $\overline{X}$ and the number of components of X by $\#X$. We will use $N(X; Y)$ to denote a closed regular neighborhood of X in Y. If the ambient space Y is clear from the context, we denote it briefly by $N(X)$. By a 3-manifold we always mean a connected and compact one, unless otherwise mentioned. Let M be a 3-manifold. Let $L \subset M$ be a submanifold with or without boundary. When L is of 1 or 2-dimension, we write $E(L) = M \setminus \operatorname{Int} N(L)$. When L is of 3-dimension, we write $E(L) = M \setminus \operatorname{Int} L$. Let X be a simplicial complex. We denote by $X^{(k)}$ the k-skeleton of X.

1. Preliminaries

Let $N_1, N_2, \ldots, N_n, L$ be possibly empty subspaces of a compact orientable 3-manifold M. We will denote by $\operatorname{Aut}(M, N_1, N_2, \ldots, N_n \text{ rel } L)$ the space of automorphisms of M which maps N_i onto N_i for any $i = 1, 2, \ldots, n$ and which is identity on L. The *mapping class group*, denoted by $\mathcal{MCG}(M, N_1, N_2, \ldots, N_n \text{ rel } L)$, is defined to be the group of isotopy classes of $\operatorname{Aut}(M, N_1, N_2, \ldots, N_n \text{ rel } L)$. Given an element f of the group of automorphisms $\operatorname{Aut}(M, N_1, N_2, \ldots, N_n \text{ rel } L)$, we use the same letter f for the representing isotopy class by abuse of notation. The "plus" subscripts, for instance in $\operatorname{Aut}_+(M, N_1, N_2, \ldots, N_n \text{ rel } L)$ and $\mathcal{MCG}_+(M, N_1, N_2, \ldots, N_n \text{ rel } L)$, indicate the subgroups of $\operatorname{Aut}(M, N_1, N_2, \ldots, N_n \text{ rel } L)$ and $\mathcal{MCG}(M, N_1, N_2, \ldots, N_n \text{ rel } L)$, respectively, consisting of orientation-preserving automorphisms (or their classes) of M.

Let $N \subset M$ be a compact 3-manifold with boundary embedded in a compact orientable 3-manifold M. An essential disk D of N is said to be *primitive* if there exists an essential disk E in $E(N)$ such that ∂D and ∂E have a single transverse intersection in ∂N. A disjoint pair of primitive disks of N is called a *primitive pair* of N.

1.1. **Handlebody-knots.** We first review the following well-known fact on the groups.

Lemma 1.1. *Let* $1 \to G' \to G \to G'' \to 1$ *be an exact sequence of groups. If* G' *and* G'' *are finitely presented, so is* G*. In particular, for a possibly empty subspaces* $N_1, N_2, \ldots, N_n$ *of a compact orientable* 3*-manifold* M*, the group* $\mathrm{Aut}(M, N_1, N_2, \ldots, N_n)$ *is finitely presented if and only if so is* $\mathrm{Aut}_+(M, N_1, N_2, \ldots, N_n)$.

Proof. The first assertion is well-known, and the particular case is also clear from the existence of following exact sequence:

$$\begin{aligned} 1 &\to \mathrm{Aut}_+(M, N_1, N_2, \ldots, N_n) \to \mathrm{Aut}(M, N_1, N_2, \ldots, N_n) && (1) \\ &\to \mathrm{Aut}(M, N_1, N_2, \ldots, N_n)/\mathrm{Aut}_+(M, N_1, N_2, \ldots, N_n) \to 1 && (2) \end{aligned}$$

and from the fact that subgroups of finite index in finitely presented groups are finitely presented, see e.g. [29]. □

A *handlebody-knot* is a handlebody embedded in the 3-sphere S^3. The *genus* of a handlebody-knot is defined to be the genus of the handlebody. A handlebody-knot V is said to be *trivial* if the exterior $E(V)$ is also a handlebody. A 2-sphere P in S^3 is called a *reducing sphere* for a handlebody-knot V if P intersects in a single essential disk. Note that $V \cap P$ and $E(V) \cap P$ are essential separating disks in V and $E(V)$ respectively. A handlebody-knot V is said to be *reducible* if there exists a reducing sphere for V. Otherwise, V is said to be *irreducible*. It is proved in [2] that a handlebody-knot V of genus two is irreducible if and only if its exterior $E(V)$ is ∂-irreducible, i.e. $\partial E(V)$ is incompressible in $E(V)$. See e.g. [18, 19] and the references given there for more details.

1.2. **Relation between the groups $\mathcal{MCG}_+(S^3, V)$ and $\mathcal{MCG}_+(E(V))$.** Let $V \subset S^3$ be a handlebody-knot.

Lemma 1.2. *The natural homomorphism* $\mathcal{MCG}_+(S^3, V) \to \mathcal{MCG}_+(E(V))$ *which takes* $f \in \mathcal{MCG}_+(S^3, V)$ *to* $f|_{E(V)} \in \mathcal{MCG}_+(E(V))$ *is an injection.*

Proof. The result follows immediately from the fact that any homeomorphism fixed on the boundary of a handlebody onto itself is isotopic to the identity (see e.g. Theorem 3.7 in [11]). □

The mapping class group $\mathcal{MCG}_+(E(V))$ is finitely presented due to the following theorem.

Theorem 1.3 ([23])**.** *Mapping class groups of irreducible orientable sufficiently large* 3*-manifolds* (*possibly not* ∂*-irreducible*) *are finitely presented.*

Following this fact, Lemma 1.2 implies that the symmetry group $\mathcal{MCG}_+(S^3, V)$ can be regarded as a subgroup of the finitely presented group $\mathcal{MCG}_+(E(V))$. However, a subgroup of a finitely generated (presented, respectively) group is not always finitely generated (presented, respectively).

1.3. **Symmetry groups of knots.**

Let K be a knot in the 3-sphere S^3. Then the group $\mathcal{MCG}(S^3, K)$ has been called the *symmetry group* [21] of the knot K and often denoted by $\mathrm{Sym}(S^3, K)$. By the uniqueness of regular neighborhoods, we have $\mathrm{Sym}(S^3, K) \cong \mathcal{MCG}(S^3, N(K))$.

The following is a direct consequence of a celebrated result of Gordon and Luecke [14] that if two knots in S^3 have homeomorphic complements then the homeomorphism between the two complements extends to an automorphism of S^3, see e.g. [21].

Proposition 1.4. *For a non-trivial knot K in S^3, we have the following natural isomorphisms:*

$$\mathcal{MCG}(S^3, N(K)) \cong \mathcal{MCG}(E(K)), \ \mathcal{MCG}_+(S^3, N(K)) \cong \mathcal{MCG}_+(E(K)).$$

By Proposition 1.4 and Theorem 1.3, we have immediately the following (the case of the unknot is clear):

Corollary 1.5. *For a knot K in S^3, the symmetry group $\mathcal{MCG}_+(S^3, N(K))$ is finitely presented.*

1.4. **Goeritz group of the trivial handlebody-knot of genus two.**

Let $V \subset S^3$ be the trivial handlebody-knot of genus g. Then the group $\mathcal{MCG}_+(S^3, V)$ is called *Goeritz group* of genus g. It was shown by Goeritz [12] in 1933 that $\mathcal{MCG}_+(S^3, V)$ is finitely generated in the case of $g = 2$. Scharlemann [26] gave a modern and concise proof of this result, and Akbas [1] and Cho [5] independently gave a finite presentation of $\mathcal{MCG}_+(S^3, V)$. We put these results together in the following theorem.

Theorem 1.6 ([12, 26, 1, 5])**.** *In the case where V is the trivial handlebody-knot of genus two, The mapping class group $\mathcal{MCG}_+(S^3, V)$ is finitely presented.*

As was mentioned in Introduction, however, the existence of such generating sets still remains unknown in the case of arbitrary genus.

2. Symmetry groups of spatial graphs

By a *graph* we shall mean a finite graph, possibly with repeated edges and *loops*, i.e. edges that connect vertices to themselves. We consider a graph as a topological space in the usual way, and hence we always assume that Γ does not have vertices of degree two unless otherwise stated. A *spatial graph* is a graph embedded in the 3-sphere S^3.

Let Γ be a spatial graph. We call the group $\mathcal{MCG}(S^3, \Gamma)$ the *symmetry group* of Γ. Note that the subgroup of the automorphism group of Γ that is induced by homeomorphisms of the pair (S^3, Γ) is called the *topological symmetry group* of the spatial graph Γ and often denoted by $\mathrm{TSG}(S^3, \Gamma)$. Such a group was introduced by Simon [28]. See [9] for details. Obviously, the group $\mathrm{TSG}(S^3, \Gamma)$ is a finite group and therefore it is always finitely presented.

The following proposition is straightforward by definition.

Lemma 2.1. *The group $\mathcal{MCG}(S^3 \text{ rel } \Gamma)$ is a normal subgroup of the symmetry group $\mathcal{MCG}(S^3, \Gamma)$ and $\mathrm{TSG}(S^3, \Gamma) \cong \mathcal{MCG}(S^3, \Gamma)/\mathcal{MCG}(S^3 \text{ rel } \Gamma)$. Hence $\mathcal{MCG}(S^3, \Gamma) \cong \mathrm{TSG}(S^3, \Gamma)$ if and only if $\mathcal{MCG}(S^3 \text{ rel } \Gamma) \cong 1$.*

2.1. Relation between symmetry groups of spatial graphs and handlebody-knots.

The proof of the following easy lemma is left to the reader.

Lemma 2.2. *Let* Γ *be a spatial graph in* S^3*. Let* f *and* g *be elements of* $\mathrm{Aut}(S^3, \Gamma)$ *such that* f *and* g *are ambient isotopic preserving* Γ *and* $f(N(\Gamma)) = g(N(\Gamma)) = N(\Gamma)$*. Then* f *and* g *determine the same element in* $\mathcal{MCG}(S^3, N(\Gamma), \Gamma)$*.*

Let V be a handlebody and let $D_1, D_2, \ldots, D_n$ be mutually disjoint, mutually non-parallel essential disks in V. Suppose that $V \setminus (\bigcup_{i=1}^n \mathrm{Int}N(D_i; V))$ consists of 3-balls. Then there exists a graph Γ embedded in $\mathrm{Int}V$ such that

(1) Γ is a deformation retract of V;
(2) Γ intersects $\bigcup_{i=1}^n D_i$ transversely at the points in the interior of the edges of Γ; and
(3) Γ has exactly n edges $e_1, e_2, \ldots, e_n$ and $\#(e_i \cap D_j) = \delta_{ij}$, where δ_{ij} is the Kronecker delta.

We call the above graph Γ the *dual graph* of $\{D_1, D_2, \ldots, D_n\}$ Note that such a graph is uniquely determined up to isotopy.

Lemma 2.3. *Let* V *be a handlebody and* Γ *be a graph embedded in* V *such that* Γ *is a deformation retract of* V*. Then the group* $\mathcal{MCG}(V, \Gamma \text{ rel } \partial V)$ *is trivial.*

Proof. Let $\{D_1, D_2, \ldots, D_n\}$ be the family of mutually disjoint, mutually non-parallel essential disks in V such that Γ is the dual graph of $\{D_1, D_2, \ldots, D_n\}$. Let $f \in \mathrm{Aut}(V, \Gamma \text{ rel } \partial V)$. Since V is irreducible, we may isotope f so that $f(D_i) = D_i$ for $1 \leqslant i \leqslant n$. Moreover, f may be isotoped so that $f|_{D_i} = \mathrm{id}_{D_i}$ by Alexander's trick. Now, the lemma follows from Alexander's trick for the complementary balls of $\bigcup_{i=1}^n D_i$. □

Let Γ be a spatial graph in S^3. Then Lemma 2.2 allows us to define in a natural way a homomorphism $\mathcal{MCG}(S^3, \Gamma) \to \mathcal{MCG}(S^3, N(\Gamma))$.

Theorem 2.4. *The above homomorphism* $\mathcal{MCG}(S^3, \Gamma) \to \mathcal{MCG}(S^3, N(\Gamma))$ *is an injection.*

Proof. Let f be an element of $\mathrm{Aut}(S^3, \Gamma)$ such that $f(N(\Gamma)) = N(\Gamma)$. Assume that f is isotopic to identity preserving $N(\Gamma)$. Then we may isotope f preserving Γ so that $f|_{E(\Gamma)} = \mathrm{id}_{E(\Gamma)}$, and thus $f|_{N(V)} \in MCG(N(\Gamma), \Gamma \text{ rel } \partial N(\Gamma))$. Now f is isotopic to the identity fixing Γ by Lemma 2.3. □

Theorem 2.4 implies that the symmetry group of a spatial graph can be regarded as a subgroup of that of a regular neighborhood of it.

2.2. Finite presentation of the symmetry groups of spatial graphs.

We review the notion of ∂-pattern in [10] and [23]. Let M be a compact 3-manifold. A *∂-pattern* for M consists of a set $\underline{\underline{m}}$ of compact connected surfaces in ∂M, such that the intersection of any i of them is a possibly empty $(n-i)$-manifold. A ∂-pattern is said to be *complete* when $\bigcup_{A \in \underline{\underline{m}}} A = \partial M$.

A ∂-pattern $\underline{\underline{m}}$ of a 3-manifold M is said to be *useful* if any disk D properly embedded in M, where ∂D intersects ∂A transversely for each $A \in \underline{\underline{m}}$ and $\#(D \cap (\bigcup_{A \in \underline{\underline{m}}} \partial A)) \leqslant 3$, bounds a disk E in ∂M such that $D \cap (\bigcup_{A \in \underline{\underline{m}}} \partial A)$ is the cone on $\partial D \cap (\bigcup_{A \in \underline{\underline{m}}} \partial A)$. An *essential* disk in $(M, \underline{\underline{m}})$ is a disk D embedded in M such that $D \cap (\bigcup_{A \in \underline{\underline{m}}} \partial A) = \emptyset$.

The mapping class group of a manifold M with ∂-pattern $\underline{\underline{m}}$, denoted by $\mathcal{MCG}(M, \underline{\underline{m}})$, is defined by $\mathcal{MCG}(M, \underline{\underline{m}}) = \mathcal{MCG}(M, A_1, A_2, \ldots, A_k)$, where $\underline{\underline{m}} = \{A_1, A_2, \ldots, A_k\}$.

Let $(M, \underline{\underline{m}})$ be a compact 3-manifold with complete ∂-pattern. *Dots* of $(M, \underline{\underline{m}})$ are pairwise disjoint points $p_1, p_2, \ldots, p_k$ in ∂M such that $p_i \cap (\bigcup_{A \in \underline{\underline{m}}} \partial A) = \emptyset$. *Spots* of $(M, \underline{\underline{m}})$ are pairwise disjoint disks $D_1, D_2, \ldots, D_k$ in ∂M such that $D_i \cap (\bigcup_{A \in \underline{\underline{m}}} \partial A) = \emptyset$.

Let $\underline{\underline{m}} = \{A_1, A_2, \ldots, A_k\}$ be a ∂-pattern of a compact 3-manifold M. The mapping class group of a dotted 3-manifold with ∂-pattern $(M, \underline{\underline{m}}, p_1, p_2, \ldots, p_k)$, denoted by $\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2, \ldots, p_k)$, is defined by

$$\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2, \ldots, p_k) = \mathcal{MCG}(M, A_1, A_2, \ldots, A_k, p_1, p_2, \ldots, p_k).$$

The mapping class group of a spotted 3-manifold with ∂-pattern $(M, \underline{\underline{m}}, D_1, D_2, \ldots, D_k)$, denoted by $\mathcal{MCG}(M, \underline{\underline{m}} \text{ rel } (D_1 \cup D_2 \cup \cdots \cup D_k)$, is defined by

$$\mathcal{MCG}(M, \underline{\underline{m}} \text{ rel } (D_1 \cup D_2 \cup \cdots \cup D_k)) = \mathcal{MCG}(M, A_1, A_2, \ldots, A_k \text{ rel } (D_1 \cup D_2 \cup \cdots \cup D_k)).$$

Lemma 2.5. *Let $(M, \underline{\underline{m}})$ be a compact 3-manifold with ∂-pattern. Assume that ∂M is connected and ∂M is neither S^2 nor $S^1 \times S^2$. Let p_1, p_2 be dots of $(M, \underline{\underline{m}})$ contained in planar components A_1 and A_2 of $\underline{\underline{m}}$ such that each component of $\partial M \setminus A_i$ is not contractible for $i = 1, 2$. Set $D_i = N(p_i; A_i)$ for $i = 1, 2$. Suppose that $\mathcal{MCG}(M, \underline{\underline{m}})$ is finitely presented. Then all of $\mathcal{MCG}(M, \underline{\underline{m}}, p_1)$, $\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$, $\mathcal{MCG}(M, \underline{\underline{m}} \text{ rel } D_1)$ and $\mathcal{MCG}(M, \underline{\underline{m}}, \text{rel}(D_1 \cup D_2))$ are finitely presented.*

Proof. Fix a set of generators $\{\gamma_1, \gamma_2, \ldots, \gamma_g\}$ of the group $\pi_1(A_1, p_1)$. Let $P_1 : \pi_1(A_1, p_1) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ be a *point-pushing map.* That is, let $\gamma : [0, 1] \to S$ be a loop in A_1 based at p_1. There is an isotopy $\varphi_t : M \to M$ supported in a small neighborhood of the loop γ such that $\varphi_0 = \text{id}$, and $\varphi_t(p_1) = \gamma(t)$. The map P_1 is then defined by $P_1(\gamma) = \varphi_1$. Since ∂M is connected and ∂M is neither S^2 nor $S^1 \times S^2$, ∂M does not admit a non-trivial S^1 action. Therefore, as a variation of the Birman exact sequence [3, 15], we have the following exact sequence:

$$1 \to \pi_1(A_1, p_1) \xrightarrow{P_1} \mathcal{MCG}(M, \underline{\underline{m}}, p_1) \to \mathcal{MCG}(M, \underline{\underline{m}}) \to 1.$$

This sequence proves that $\mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ is finitely presented by Lemma 1.1.

If $A_1 \neq A_2$, then it is clear from the above argument that $\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$ is finitely presented. Let $A_1 = A_2 = A$ and $P_2 : \pi_1(A \setminus \{p_1\}, p_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$ be a point-pushing map. Then the Birman exact sequence

$$1 \to \pi_1(A \setminus \{p_1\}, p_2) \xrightarrow{P_2} \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1) \to 1,$$

shows that $\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$ is finitely presented.

Let $\tau_1 \in \mathcal{MCG}(M, \underline{\underline{m}}, D_1)$ be the Dehn twist about D_1. Then we have the following exact sequence:

$$1 \to \langle \tau_1 \rangle \to \mathcal{MCG}(M, \underline{\underline{m}}, D_1) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1) \to 1,$$

where the map $\mathcal{MCG}(M, \underline{\underline{m}}, D_1) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ is induced by collapsing the spot D_1 to p_1. This implies that $\mathcal{MCG}(M, \underline{\underline{m}}, D_1)$ is finitely presented.

Finally, let $\tau'_i \in \mathcal{MCG}(M, \underline{\underline{m}}, D_1, D_2)$ be the Dehn twist about D_i for $i = 1, 2$. Then considering the following exact sequence:

$$1 \to \langle \tau_1, \tau_2 \rangle \to \mathcal{MCG}(M, \underline{\underline{m}}, D_1, D_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2) \to 1,$$

it follows that $\mathcal{MCG}(M, \underline{\underline{m}}, D_1)$ is finitely presented. □

Lemma 2.6. *Let* $(M, \underline{\underline{m}})$ *be a compact* 3*-manifold with* ∂*-pattern such that* $\partial M \cong S^1 \times S^1$*. Let* p_1, p_2 *be dots of* $(M, \underline{\underline{m}})$ *contained in components* A_1 *and* A_2 *of* $\underline{\underline{m}}$ *such that* $A_i \cong D^2$ *or* $A_i \cong S^1 \times [0, 1]$ *and the inclusion* $A_i \hookrightarrow \partial M$ *induces an injection of their fundamental groups for* $i = 1, 2$*. Set* $D_i = N(p_i; A)$ *for* $i = 1, 2$*. Suppose that* $\mathcal{MCG}(M, \underline{\underline{m}})$ *is finitely presented. Then all of* $\mathcal{MCG}(M, \underline{\underline{m}}, p_1)$*,* $\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$*,* $\mathcal{MCG}(M, \underline{\underline{m}} \text{ rel } D_1)$ *and* $\mathcal{MCG}(M, \underline{\underline{m}} \text{ rel } (D_1 \cup D_2))$ *are finitely presented.*

Proof. We give the proof for the case $A_1 = A_2 = A$; the other case is left to the reader.

Suppose that A is a disk. Then we immediately have an isomorphism

$$\mathcal{MCG}(M, \underline{\underline{m}}, p_1) \cong \mathcal{MCG}(M, \underline{\underline{m}}).$$

It follows from Lemma 1.1 that $\mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ is finitely generated.

We fix a single generator γ of the group $\pi_1(A \setminus \{p_1\}, p_2)$. Let $P_2 : \pi_1(A \setminus \{p_1\}, p_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$ be a point-pushing map. The map P_2 induces a map $\bar{P}_2 : \pi_1(A \setminus \{p_1\}, p_2)/\ker P_2 \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$. Then there exists the following exact sequence

$$1 \to \pi_1(A \setminus \{p_1\}, p_2)/\ker P_2 \xrightarrow{\bar{P}} \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2) \to \mathcal{MCG}(M, \underline{\underline{m}}) \to 1.$$

Since $\pi_1(A \setminus \{p_1\}, p_2) = \langle \gamma \rangle \cong \mathbb{Z}$, the group $\pi_1(A \setminus \{p_1\}, p_2)/\ker P_2$ is finitely presented. Hence $\mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ is finitely generated by Lemma 1.1.

Next, suppose that A is an annulus. Fix a unique generator γ the group $\pi_1(A, p_1)$. Let $P'_1 : \pi_1(A, p_1) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ be a point-pushing map. The map P'_1 induces a map $\bar{P}'_1 : \pi_1(A, p_1)/\ker P'_1 \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1)$. Then there exists the Birman exact sequence

$$1 \to \pi_1(A \setminus \{p_1\}, p_2)/\ker P'_1 \xrightarrow{\bar{P}'_1} \mathcal{MCG}(M, \underline{\underline{m}}, p_1) \to \mathcal{MCG}(M, \underline{\underline{m}}) \to 1.$$

Since $\pi_1(A, p_1) = \langle \gamma \rangle \cong \mathbb{Z}$, the group $\pi_1(A, p_1)/\ker P'_1$ is finitely presented. Hence $\mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ is finitely generated by Lemma 1.1.

Let $P'_2 : \pi_1(A \setminus \{p_1\}, p_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$ be a point-pushing map. Then by the Birman exact sequence

$$1 \to \pi_1(A \setminus \{p_1\}, p_2) \xrightarrow{P'_2} \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1) \to 1$$

it follows that $\mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2)$ is finitely generated by Lemma 1.1.

Let $\tau_1 \in \mathcal{MCG}(M, \underline{\underline{m}}, D_1)$ be the Dehn twist about D_1. Then we have the following exact sequence:

$$1 \to \langle \tau_1 \rangle \to \mathcal{MCG}(M, \underline{\underline{m}}, D_1) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1) \to 1,$$

where the map $\mathcal{MCG}(M, \underline{\underline{m}}, D_1) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1)$ is induced by collapsing the spot D_1 to p_1. This implies that $\mathcal{MCG}(M, \underline{\underline{m}}, D_1)$ is finitely presented.

Finally, let $\tau_i' \in \mathcal{MCG}(M, \underline{\underline{m}}, D_1, D_2)$ be the Dehn twist about D_i for $i = 1, 2$. Then considering the following exact sequence:

$$1 \to \langle \tau_1, \tau_2 \rangle \to \mathcal{MCG}(M, \underline{\underline{m}}, D_1, D_2) \to \mathcal{MCG}(M, \underline{\underline{m}}, p_1, p_2) \to 1,$$

it follows that $\mathcal{MCG}(M, \underline{\underline{m}}, D_1)$ is finitely presented. □

Let K be a knot in S^3. Let D be a meridian disk of $N(K)$. Set $A = \partial N(K) \cap N(D; N(K))$ Regard A as $S^1 \times [0, 1]$, and divide it into three annuli $A_1 = S^1 \times [0, 1/3]$, $A_2 = S^1 \times [1/3, 2/3]$ and $A_3 = S^1 \times [2/3, 1]$. Let $B = \partial N(K) \setminus \mathrm{Int} A$. Then the complete ∂-pattern $\underline{\underline{m}} = \{A_1, A_2, A_3, B\}$ satisfies the following:

Lemma 2.7. *$\mathcal{MCG}(E(K), \underline{\underline{m}})$ is finitely presented.*

Let Γ be a spatial graph in S^3. Set $V = N(\Gamma)$. Let $\{D_1, e_2, \ldots, D_n\}$ be the family of essential disks in V such that Γ is the dual graph of $\{D_1, e_2, \ldots, D_n\}$. Set $A_i = \partial V \cap N(D_i; V)$ for $1 \leqslant i \leqslant n$. Let $B_1, B_2, \ldots, B_m$ be the set of components of $\partial V \setminus \bigcup_{i=1}^n \mathrm{Int}\, A_i$. Then $\underline{\underline{m}} = \{A_i\}_{i=1}^n \cup \{B_j\}_{j=1}^m$ is a complete ∂-pattern of $E(\Gamma) = E(V)$.

For each essential disk D_i in $N(\Gamma)$ corresponding to a loop e_i of Γ, regard $N(D_i; N(\Gamma)) \cap \partial N(\Gamma)$ as $S^1 \times [0, 1]$, and divide it into three annuli $S^1 \times [0, 1/3]$, $S^1 \times [1/3, 2/3]$ and $S^1 \times [2/3, 1]$. Applying this operation for every loop of Γ, we obtain a new ∂-pattern $\underline{\underline{n}}$ of $E(\Gamma)$.

Lemma 2.8. *$\mathcal{MCG}(E(\Gamma), \underline{\underline{n}})$ is finitely presented.*

The key lemma for the proof of the above lemma is the following theorem in [4]:

Theorem 2.9 ([4]). *Suppose a group G acts cellularly on a contractible CW-complex $\mathcal{K}$. If the 2-skeleton of $\mathcal{T}/G$ is finite, the stabilizer of each 0-cell of $\mathcal{K}$ is finitely presented, and the stabilizer of each 1-cell of $\mathcal{K}$ is finitely generated, then G is finitely presented.*

Proof of Lemma 2.8. Assume that $E(\Gamma)$ is ∂-irreducible. Then $E(\Gamma)$ is a compact, irreducible, sufficiently large 3-manifold and $\underline{\underline{n}}$ is a useful ∂-pattern of $E(\Gamma)$. Hence it is straightforward from [23] that $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}})$ is finitely presented.

We now turn to the case where $\partial E(\Gamma)$ is compressible in $E(\Gamma)$. In the following, we include the case where Γ consists of a single vertex and a single loop. If $\underline{\underline{n}}$ is still a useful ∂-pattern of $E(\Gamma)$, then we are done. Assume that there exists a disk D properly embedded in $E(\Gamma)$ such that ∂D intersects ∂A transversely for each $A \in \underline{\underline{n}}$ and $\#(D \cap \bigcup_{A \in \underline{\underline{n}}} \partial A) \leqslant 3$. Then it is clear from the definition of $\underline{\underline{n}}$ that $\#(D \cap \bigcup_{A \in \underline{\underline{m}}} \partial A) = 0$.

The proof is an induction of the genus of $N(\Gamma)$. We proved in Lemma 2.7 the case where the genus of $N(\Gamma)$ is one. Assume that the assertion holds if the genus of $N(\Gamma)$ is less than g and Let $N(\Gamma) = g$.

We denote by $\mathcal{Z}$ the simplicial complex such that

1. the set of vertices of $\mathcal{Z}$ consists of the ambient isotopy classes (in $(E(\Gamma))$ of essential disks in $(E(\Gamma), \underline{\underline{n}})$;
2. a collection of $k + 1$ vertices spans a k-simplex if and only if they admit a set of pairwise-disjoint representatives.

Note that $\mathcal{Z}$ is a subcomplex of the *disk complex* of $E(\Gamma)$, see [23] for the definition of disk complexes. Let E_1 and E_2 be distinct essential disks in $(E(\Gamma), \underline{\underline{n}})$. Assume that $\#(E_1 \cap E_2)$ is minimal up to isotopy. Suppose that $E_1 \cap E_2 \neq \emptyset$. Let $A \subset E_1$ be a disk cut off from E_1 by an outermost arc α of $E_1 \cap E_2$ in E_1 such that $A \cap E_2 = \emptyset$. Then both disks obtained from surgery on E_2 along A are vertices of $\mathcal{Z}$. It follows from the proof of Theorem 4.2 in [5] that $\mathcal{Z}$ is contractible.

Let $\{E_1, E_2\}$ be an edge of $\mathcal{Z}$. Then there exist elements A_1, A_2 of $\underline{\underline{n}}$ (possibly $A_1 = A_2$) such that $\partial E_1 \subset A_1$ and $\partial E_2 \subset A_2$. Note that we may assume that neither A_1 nor A_2 is an annulus. The simple closed curves ∂E_1 and ∂E_2 bound disks $\hat{E}_1$, $\hat{E}_2$ in $N(\Gamma)$, respectively. Let $\Gamma_{\{E_1,E_2\}}$ be the graph embedded in $N(\Gamma)$ such that $\Gamma_{\{E_1,E_2\}}$ is the dual of $\{D_1, D_2, \ldots, D_n, \hat{E}_1, \hat{E}_2\}$. Here, if $\hat{E}_i$ is parallel to a disk D_j of $\{D_1, D_2, \ldots, D_n\}$ then we remove D_j from the family $\{D_1, D_2, \ldots, D_n, \hat{E}_1, \hat{E}_2\}$. Let e'_1 and e'_2 be the edges of $\Gamma_{\{E_1,E_2\}}$ corresponding to the disks $\hat{E}_1$ and $\hat{E}_2$. Then the labeled spatial graph $(\Gamma_{\{E_1,E_2\}}, e'_1, e'_2)$ is obtained from Γ by one of the following local operations:

1. Choose arbitrary two (cut) edges from Γ and label them by e'_1 and e'_2;
2. Choose an arbitrary (cut) edge of Γ and label it by e'_1 (e'_2, respectively). Let v be a degree k (cut) vertex of Γ and let C be a disk properly embedded in $N(v; S^3)$ such that $C \cap \Gamma = N(v; S^3) \cap \Gamma$ and that $C \cap \Gamma$ is a cone on k points. Replace $\Gamma \cap C$ by the tree on C whose degree one edges are exactly $\Gamma \cap \partial C$ and the other two vertices v_1 and v_2 are such that
   (a) $\deg(v_1) > 2$ and $\deg(v_2) > 2$; and
   (b) $\deg(v_1) + \deg(v_2) = k + 2$.
   Label the edge connecting v_1 and v_2 by e'_2 (e'_1, respectively), see Figure 1;



FIGURE 1. From Γ to $(\Gamma_{\{E_1,E_2\}}, e'_1, e'_2)$.

3. Let v be a degree k (cut) vertex of Γ and let C be a disk properly embedded in $N(v; S^3)$ such that $C \cap \Gamma = N(v; S^3) \cap \Gamma$ and that $C \cap \Gamma$ is a cone on k points. Replace $\Gamma \cap C$ by the tree on C whose degree one edges are exactly $\Gamma \cap \partial C$ and the other four vertices v', v_1, v_2 and v_3 are such that
   (a) $\deg(v) = 3$;
   (b) v is incident to v_1, v_2 and v_3; and
   (c) $\deg(v_2) > 2$, $\deg(v_2) > 2$ and , $\deg(v_3) > 2$; and
   (d) $\deg(v_1) + \deg(v_2) + \deg(v_3) = k + 4$.
   Label the edge connecting v' and v_1 by e'_1 and the edge connecting v' and v_2 by e'_2, see Figure 2;
4. Let v and v' be (cut) vertices of Γ such that $\deg(v) = k$ and $\deg(v') = k'$. Let C (C', respectively) be a disk properly embedded in $N(v; S^3)$ ($N(v'; S^3)$,



FIGURE 2. From Γ to $(\Gamma_{\{E_1,E_2\}}, e'_1, e'_2)$.

respectively) such that $C \cap \Gamma = N(v; S^3) \cap \Gamma$ ($C' \cap \Gamma = N(v'; S^3) \cap \Gamma$, respectively) and that $C \cap \Gamma$ ($C' \cap \Gamma$, respectively) is a cone on k (k', respectively) points. Replace $\Gamma \cap C$ ($\Gamma \cap C'$, respectively) by the tree on C whose degree one edges are exactly $\Gamma \cap \partial C$ and the other two vertices v_1 and v_2 (v'_1 and v'_2, respectively) are such that

(a) $\deg(v_1) > 2$ and $\deg(v_2) > 2$ ($\deg(v'_1) > 2$ and $\deg(v'_2) > 2$, respectively); and
(b) $\deg(v_1) + \deg(v_2) = k + 2$ ($\deg(v'_1) + \deg(v'_2) = k' + 2$, respectively).

Label the edge connecting v_1 and v_2 by e'_1 and the edge connecting v'_1 and v'_2 by e'_2.

Since each of above operations may only produce finitely many labeled spatial graphs $\Gamma_{\{E'_1,E'_2\}}$, it follows that the 2-skeleton of $\mathcal{Z}/\mathcal{MCG}(E(\Gamma), \underline{\underline{n}})$ is finite.

Let E be a vertex of $\mathcal{Z}$. We proceed to show that $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}}, E)$ is finitely presented. Cutting $E(\Gamma)$ along E, we get two manifolds L_1 and L_2. Both ∂L_1 and ∂L_2 inherit ∂-patterns $\underline{\underline{l_1}}$ and $\underline{\underline{l_2}}$ from $\underline{\underline{m}}$. Let E' and E'' be cut end disks on ∂L_1 and ∂L_2, respectively. Note that E' and E'' are spots of $(L_1, \underline{\underline{l_1}})$ and $(L_2, \underline{\underline{l_2}})$, respectively. By the assumption of the induction and Lemmas 2.5 and 2.6, it follows that both $\mathcal{MCG}(L_1, \underline{\underline{l_1}} \text{ rel } E')$ and $\mathcal{MCG}(L_2, \underline{\underline{l_2}} \text{ rel } E'')$ are finitely presented.

Let $\tau_1 \in \mathcal{MCG}_+(L_1, \underline{\underline{l_1}} \text{ rel } E')$ and $\tau_2 \in \mathcal{MCG}_+(L_2, \underline{\underline{l_2}} \text{ rel } E'')$ be respectively Dehn twists about E' and E''. Then we have

$$\mathcal{MCG}_+(E(\Gamma), \underline{\underline{n}}, E) \cong \mathcal{MCG}_+(L_1, \underline{\underline{l_1}} \text{ rel } E') *_{\langle \tau_1 \tau_2^{-1} \rangle} \mathcal{MCG}_+(L_2, \underline{\underline{l_2}} \text{ rel } E'')$$

if there is no element in $\mathcal{MCG}_+(E(\Gamma), \underline{\underline{n}}, E)$ which restricts to an orientation reversing automorphism of E and

$$\mathcal{MCG}_+(E(\Gamma), \underline{\underline{n}}, E) \cong \mathcal{MCG}_+(L_1, \underline{\underline{l_1}} \text{ rel } E') *_{\langle \tau_1 \tau_2^{-1} \rangle} \mathcal{MCG}_+(L_2, \underline{\underline{l_2}} \text{ rel } E'') \rtimes \mathbb{Z}/2\mathbb{Z}$$

otherwise. Now it follows from Lemma 1.1 that $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}}, E)$ is finitely presented.

Let $\{E_1 E_2\}$ be an edge of $\mathcal{Z}$. Then we may prove that $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}}, E_1 \cup E_2)$ is finitely generated in an analogous argument using Lemmas 2.5 and 2.6 as above.

Consequently, the action $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}})$ on the contractible simplicial complex $\mathcal{Z}$ satisfies the assumption of Theorem 2.9 and thus we get that $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}})$ is finitely presented, which is our claim.

Theorem 2.10. *For a spatial graph Γ in S^3, the symmetry group $\mathcal{MCG}(S^3, \Gamma)$ is finitely presented.*

Proof. It is clear that $\mathcal{MCG}(E(\Gamma), \underline{\underline{n}}) \cong \mathcal{MCG}(S^3 \text{ rel } \Gamma)$. By Lemma 2.1 there exists the following exact sequence:

$$1 \to \mathcal{MCG}(S^3 \text{ rel } \Gamma) \to \mathcal{MCG}(S^3, \Gamma) \to \mathrm{TSG}(S^3, \Gamma) \to 1.$$

Since both $\mathcal{MCG}(S^3 \text{ rel } \Gamma)$ and $\mathrm{TSG}(S^3, \Gamma)$ are finitely presented due to Lemma 2.8, it follows from Lemma 1.1 that $\mathcal{MCG}(S^3, \Gamma)$ is finitely presented.

□

Example. Figure 3 illustrates a *handcuff graph* Γ planarly embedded in S^3. Note that the graph Γ has the unique edge e that is not a loop. This implies that all elements of $\mathcal{MCG}(S^3, \Gamma)$ preserve e. Thus we have



Figure 3. A planarly embedded handcuff graph.

$$\mathcal{MCG}_+(S^3, \Gamma) = \langle g_1, g_2, g_3 \mid g_3 g_1 g_3 {g_1}^{-1}, g_3 g_2 g_3 {g_2}^{-1} {g_1}^{-1} \rangle,$$

where g_1, g_2, g_3 are shown in Figure 4.



Figure 4. Automorphisms g_1, g_2 and g_3.

3. The symmetry groups of reducible handlebody-knots of genus two

Let $V \subset S^3$ be a handlebody-knot of genus two. In this section we assume that there exists a reducing sphere P for V. The sphere P separates S^3 into two 3-balls B_1 and B_2, i.e. $S^3 = B_1 \cup_P B_2$. Set $V_i = V \cap B_i$ for $i = 1, 2$ and Let K_1 and K_2 be the cores of V_1 and V_2 respectively. Set $M_i = E(V) \cap B_i$ for $i = 1, 2$. Note that M_i is homeomorphic to $E(K_i)$.

3.1. The case where both K_1 and K_2 are non-trivial.

In this subsection we always assume that both K_1 and K_2 are non-trivial knots. Figure 5 illustrates a typical picture of V in this case.

Lemma 3.1. *If both K_1 and K_2 are non-trivial, $E(V) \cap P$ is the unique* (*up to ambient isotopy*) *compression disk of $\partial E(V)$ in $E(V)$.*



Figure 5. A typical case.

Proof. Set $D = E(V) \cap P$. Let D_1 be an essential disk in $E(V)$. Isotope D_1 in order to minimize transverse intersections of $D \cap D_1$ and assume, in contradiction, that $D \cap D_1 \neq \emptyset$. Since $E(V)$ is irreducible, there exists not loop in $D \cap D_1$. let $A \subset D_1$ be a disk cut off from D_1 by an outermost arc α of $D \cap D_1$ in D_1 such that $A \cap D = \alpha$. We may assume that A is contained in M_1. Then A is a proper disk in $M_1 \cong E(K_1)$, and since K_1 is assumed to be non-trivial, ∂A bounds a disk A' in ∂M_1. Set $\beta = A' \cap \partial D$. Since $E(V)$ is irreducible, $A \cap A'$ bounds a 3-ball B in $E(V)$. Therefore we may isotopy D_1 in $N(B)$ until it is disjoint from D, and this isotopy decreases $\#(D \cap D_1)$, contradicting minimality of $\#(D \cap D_1)$. Thus $D \cap D_1 = \emptyset$. We may assume, without loss of generality, that $D_1 \subset M_1$. Since K_1 is non-trivial, $M_1 \cong E(K_1)$ is Haken. Hence ∂D_1 bounds a disk A'' in ∂M_1. By assumption, ∂D_1 is essential in $\partial E(V)$, thus essential in $\partial E(V) \cap \partial M_1$ as well. Thus A'' must contain $D \subset \partial M_1$. We conclude, by the irreducibility of $E(V)$, that D_1 is parallel to D. □

Remark that from the above Lemma and irreducibility of the handlebody V, it turns out that P is the unique reducing sphere for V.

Lemma 3.2. *If both K_1 and K_2 are non-trivial, we have the natural isomorphism $\mathcal{MCG}_+(S^3, V) \cong \mathcal{MCG}_+(E(V))$.*

Proof. Let $\varphi : \mathcal{MCG}_+(S^3, V) \to \mathcal{MCG}_+(E(V))$ be the natural homomorphism which takes $f \in \mathcal{MCG}_+(S^3, V)$ to $f|_{E(V)} \in \mathcal{MCG}_+(E(V))$. By Lemma 1.2, it remains to show the surjectivity of φ. Set $D = E(V) \cap P$. Take an arbitrary element $g \in \mathrm{Aut}_+(E(V))$. By Lemma 3.1, we may isotope g so that $g(D) = D$. It suffices to show that g extends to an element of $\mathrm{Aut}_+(S^3, V)$.

Assume that $g|_D$ is orientation preserving. Then we have that $g|_{M_i} \in \mathrm{Aut}_+(M_i)$. Since $M_i \cong E(K_i)$, it follows from [14] that $g|_{M_i}$ extends to an element of $\mathrm{Aut}_+(B_i, V_i)$. Thus g extends to an element of $\mathrm{Aut}_+(S^3, V)$ by Alexander's trick. The situation is only slightly more complicated if we drop the requirement that $g|_D$ be orientation preserving. Assume that $g|_D$ is orientation reversing. Note that in this case K_1 and K_2 are the same knot. Then $g(M_i) = (M_{i+1})$ (subscripts mod 2). Again due to [14], we see that g takes a meridian slope of V_i in $\partial M_i \setminus \mathrm{Int}\, D$ to a meridian slope of V_{i+1} in $\partial M_{i+1} \setminus \mathrm{Int}\, D$. Therefore g extends to an element of $\mathrm{Aut}_+(S^3, V)$. □

Now, the following proposition follows immediately from Lemmas 1.1 and 3.2, and Theorem 1.3.

Proposition 3.3. *If both K_1 and K_2 are non-trivial, then the symmetry group $\mathcal{MCG}(S^3, V)$ of V is finitely presented.*

3.2. **The case where K_1 is non-trivial and K_2 is trivial.** In this subsection we always assume that only one of them (say K_2) is the unknot. (Hence K_1 is non-trivial.) Figure 6 illustrates a typical picture of V in this case.



FIGURE 6. A typical case.

Let W be a 3-manifold obtained from a connected orientable surface F by attaching a single 2-handle to $F \times \{1\} \subset F \times [0, 1]$ along a non-separating simple closed curve l in $F \times \{1\}$. Let $\partial_+ W = F \times \{0\}$ and $\partial_- W = \partial W \setminus \partial_+ W$. The cores of the 2-handle can be vertically extended to a non-separating essential disk E through $F \times [0, 1]$.

Lemma 3.4. *Any non-separating essential disk E' in W such that $\partial E' \subset \partial_+ W$ is parallel to E.*

Proof. Let E' be a non-separating essential disk in W such that $\partial E' \subset \partial_+ W$. Isotope E' in order to minimize $\#(E \cap E')$ and assume, in contradiction, that $E \cap E' \neq \emptyset$. It is easy to see that W is irreducible, hence all components of $E \cap E'$ are arcs. Let $A \subset E'$ be a disk cut off from E' by an outermost arc α of $E \cap E'$ in E' such that $A \cap E = \alpha$. Then A is a proper disk in a 3-manifold W' obtained by cutting W along E. Since W' is homeomorphic to $\partial_- W \times [0, 1]$ and $\partial_- W \times [0, 1]$ is ∂-irreducible, ∂A bounds a disk A' in $\partial W'$. Since $\partial_- W \times [0, 1]$ is irreducible, $A \cup A'$ bounds a 3-ball in W'. This 3-ball gives an instruction about how to isotope E' until it is disjoint from E around the 3-ball. This isotopy decreases $\#(E \cap E')$, contradicting minimality of $\#(E \cap E')$. Therefore we may assume that E and E' are disjoint. Now, the lemma follows from the fact that W' is irreducible and ∂-irreducible. □

Lemma 3.5. *Any two primitive disks E and E' of $E(V)$ are parallel.*

Proof. Set $W = M_2 \cup N(\partial M_1; M_1)$. The boundary ∂W of W consists of two components $\partial_+ W = \partial E(V)$ and $\partial_- W = \partial W \setminus \partial_+ W$.

The surface $\partial_- W$ separates $E(V)$ into two components W and $E(W) = V \setminus \operatorname{Int} W$. Then it is clear that the surface $\partial_- W$ is incompressible in W since the negative boundary of a compression body is incompressible. The surface $\partial_- W$ is also incompressible in $E(W)$ since $(E(W), \partial_- W)$ is homeomorphic to $(E(K), \partial E(K))$. Thus $\partial_- W$ is incompressible in $E(V)$. Since $E(V)$ is irreducible and $\partial_- W$ is incompressible in $E(V)$, The disk E' can be isotoped into W. Thus the result follows immediately from Lemma 3.4. $\square$

Lemma 3.6. *Let H be the genus two handlebody. Let $D_2 \subset H$ be an essential non-separating disk. Let l be a non-separating simple closed curve in ∂H such that ∂D_2 and l have a single transverse intersection in ∂H. Let $H(l)$ be a compact 3-manifold obtained by adding to H a 2-handle along l. (Hence $H(l) \cong D^2 \times S^1$.) Let $D_1 \subset H$ be an essential non-separating disk such that $D_1 \cap (D_2 \cup l) = \emptyset$ and that D_1 is a meridian disk of $H(l)$. If $D' \subset H$ be an essential non-separating disk such that $\partial D'$ and l have a single transverse intersection in ∂H, D' can be isotoped in H so that $D_1 \cap D' = \emptyset$.*

Proof. Embed H in S^3 so that both D_1 is a primitive disk and l bounds a disk E in $E(H)$. Isotope D' in order to minimize transverse intersections of $D_1 \cap D'$ and assume, in contradiction, that $D_1 \cap D' \neq \emptyset$. We may assume that $\#(D_1 \cap D')$ is minimal among all such disks D'. Let $A \subset D_1$ be a disk cut off from D_1 by an outermost arc α of $D_1 \cap D'$ in D_1 such that $A \cap D' = \alpha$. Let D'_1 and D'_2 are disks obtained from surgery on D' along A. By Lemma 2.3 in [5], both D'_1 and D'_2 are primitive. We may assume that $D'_1 \cap l = \emptyset$ and $\partial D'_2$ and l have a single transverse intersection.

Claim. The disk D'_1 is parallel to D_1.

Proof of Claim. Since $\#(D_1 \cap D'_2) < \#(D_1 \cap D')$, one can isotope D_1 off of D'_2 by the minimality of $\#(D_1 \cap D')$. Recall that D'_1 is also disjoint from D'_2. Let $H' = H \cup N(E)$. Then all meridian disks of H' disjoint from $N(E) \cup D'_2$ are parallel. Since D_1 and D'_1 are primitive and we can isotope these disks off of D'_2, D_1 and D'_1 are meridian disks of H' disjoint from D'_2, and hence they are parallel.

The disk D' is reconstructed by tubing D'_1 and D'_2 along a simple arc having one endpoint on D'_1 and the other one on D'_2. This implies that D' can be isotoped so as to be disjoint from D_1, contradicting the choice of D'. $\square$

The following lemma is an immediate consequence of Lemma 3.5 and 3.6.

Lemma 3.7. *Let D_1 be a meridian disk of the solid torus V_1 such that D_1 is disjoint from the reducing sphere P. Then any primitive disk D' of V can be isotoped in V so that $D_1 \cap D' = \emptyset$.*

The *primitive disk complex* $\mathcal{P}(V)$ of $V \subset S^3$ is defined to be the simplicial complex such that

1. the set of vertices of $\mathcal{P}(V)$ consists of the ambient isotopy classes of primitive disks in V;
2. a collection of $k+1$ vertices spans a k-simplex if and only if they admit a set of pairwise-disjoint representatives.

Note that the primitive disk complex $\mathcal{P}(V)$ is a subcomplex of a *disk complex* of V.

Lemma 3.8. *The primitive disk complex $\mathcal{P}(V)$ is contractible.*

Proof. Let D_2 and D'_2 be primitive disks in V. By Lemma 3.5, we may assume that the slopes ∂D_2 and ∂E ($\partial D'_2$ and ∂E) have a single transverse intersection in ∂V, and that

$\partial D_2 \cap \partial D_2' \cap \partial E = \emptyset$. Choose a representation of D_2' of its ambient isotopy class which satisfies the above condition and which minimizes $\#(D_2 \cap D_2')$. Assume that $D_2 \cap D_2' \neq \emptyset$. Let $A \subset D_2$ be a disk cut off from D_2 by an outermost arc α of $D_2 \cap D_2'$ in D_2 such that $A \cap D_2' = \emptyset$. The boundary $\partial\alpha$ of α separates ∂D_2 ($\partial D_2'$, respectively) into two arcs δ_1 and δ_2 (δ_1' and δ_2', respectively). Let F_1 and F_2 be disks obtained from surgery on D_2' along A. We may assume that $\partial A = \delta_1 \cup \alpha$, $\delta_1' \cap l \neq \emptyset$, and $\partial F_i = \delta_1 \cup \delta_i'$ for $i = 1, 2$. If δ_1 intersects l, the slopes ∂F_2 and ∂E have a single transverse intersection in ∂V. If δ_1 does not intersect l, the slopes ∂F_1 and ∂E have a single transverse intersection in ∂V. Therefore one of F_1 and F_2 is primitive. Now the lemma follows from Theorem 4.2 in [5]. □

Lemma 3.9. *The primitive disk complex $\mathcal{P}(V)$ is of 1-dimension.*

Proof. It is clear that the dimension of $\mathcal{P}(V)$ is at least 1. Let D_2, D_2' and D_2'' be pairwise disjoint primitive disks of V. By , we may assume that the three disks do not intersect a meridian disk D_1 of the solid torus V_1, where we assume that D_1 is disjoint from the reducing sphere P. Thus D_2, D_2' and D_2'' can be regarded as meridian disks of the solid torus $V' = V \setminus \operatorname{Int} N(D_1; V)$. Cut V' along $D_2 \cup D_2' \cup D_2''$. Then V' decomposes into three components and at least one of them does not intersect $N(D_1; V)$. This implies at least two of the three disk are parallel in V. □

Lemma 3.10. *Let D be a primitive disk of V. Then the solid torus $V \setminus \operatorname{Int} N(D)$ is ambient isotopic to the solid torus $N(K_1)$.*

Proof. By Lemma 3.5), there is the unique primitive disk of $E(V)$. Let D_2 be a meridian disk of V_2 such that $\partial D_2 \cap P = \emptyset$. We may assume that ∂D_2 and ∂E have a single transverse intersection. Regard $N(E; E(V))$ as a product $E \times [0,1]$ and $N(D_2; V)$ as $D_2 \times [0,1]$. Note that $V \cup E \times [0,1]$ is ambient isotopic to $N(K_1)$ since $V \cup (E \times [0,1])$ is obtained by attaching the 3-ball $V_2 \cup (E \times [0,1])$ to $V_1 \cong N(K_1)$ along a disk $P \cap V$, $V \cup (E \times [0,1])$ is ambient isotopic to $N(K_1)$. The union $V \cup (E \times [0,1])$ is also obtained from $V \setminus (D_2 \times (0,1))$ by attaching a 3-ball $(E \times [0,1]) \cup (D_2 \times [0,1])$ along a disk $(D \times \{0,1\}) \cup ((\partial E \times [0,1]) \setminus (\partial D_2 \times (0,1)))$, we have that $V \setminus (D_2 \times (0,1))$ is ambient isotopic to $V \cup (E \times [0,1])$.

Let D_2' be an arbitrary primitive disk in V. Then we may assume, again, that $\partial D_2'$ and ∂E have a single transverse intersection due to the uniqueness of E. By the same argument as above, we have that $V \setminus \operatorname{Int} N(D_2'; V)$ is ambient isotopic to $V \cup (E \times [0,1])$. This completes the proof. □

Lemma 3.11. *The group $\mathcal{MCG}_+(S^3, V)$ acts transitively on the set of primitive disks of V. The group $\mathcal{MCG}_+(S^3, V)$ acts transitively on the set of primitive pairs of V.*

Proof. Let D_2 and D_2' be primitive disks of V. Due to Lemma 3.10, we may regard $N(D_2; V)$ and $N(D_2'; V)$ as regular neighborhoods of ∂-parallel proper arcs in $E(K_1)$. Since any two ∂ arcs in $E(K_1)$ are ambient isotopic in $E(K_1)$, we get the first assertion.

Let $\{D_2, D_2'\}$ and $\{D_3, D_3'\}$ be primitive pairs of V. We may assume that Each of these primitive disks has a single transverse intersection with ∂E. Set $p_2 = \partial D_2 \cap E$ and $q_2 = \partial D_2' \cap E$ ($p_3 = \partial D_3 \cap E$ and $q_3 = \partial D_3' \cap E$, respectively). Let α_2 (α_3, respectively) be a simple proper arc in E connecting p_2 and q_2 (p_3 and q_3, respectively).

Regard $N(E; E(V))$ as a product $E \times [0,1]$. Then the union of the arcs $(\partial D_2 \cup \partial D_2') \setminus (\partial E \times (0,1))$ ($(\partial D_3 \cup \partial D_3') \setminus (\partial E \times (0,1))$, respectively) and the arcs $\alpha_2 \times \{0,1\}$ ($\alpha_3 \times \{0,1\}$, respectively) is a simple closed curve in the boundary of the solid torus $V \cup (E \times [0,1]) \cong E(K_1)$ bounding a disk in $V \cup (E \times [0,1])$. Since any simple closed curve in $\partial E(K_1)$ bounding a disk in $E(K_1)$ are ambient isotopic, we get the second assertion.

□

Let $\mathcal{P}'(V)$ be the first barycentric subdivision of $\mathcal{P}(V)$. By Lemma 3.11, the quotient of $\mathcal{P}'(V)$ by the action of $\mathcal{MCG}_+(S^3, V)$ is a single edge. For convenience, we will not distinguish disks and homeomorphisms from their ambient isotopy classes in their notations. Fix two vertices $v_1 = D_2$ and $v_2 = \{D_2, D_2'\}$ that are endpoints of an edge e of $\mathcal{P}'(V)$. Note that $\mathrm{Stab}_{\mathcal{MCG}_+(S^3,V)}(v_1) = \mathcal{MCG}_+(S^3, V, D_2)$, $\mathrm{Stab}_{\mathcal{MCG}_+(S^3,V)}(v_2) = \mathcal{MCG}_+(S^3, V, D_2 \cup D_2')$ and $\mathrm{Stab}_{\mathcal{MCG}_+(S^3,V)}(e) = \mathcal{MCG}_+(S^3, V, D_2, D_2')$.

By the theory of groups acting on trees due to Bass and Serre [27], the mapping class group $\mathcal{MCG}_+(S^3, V)$ is a free product with amalgamation

$$\mathcal{MCG}_+(S^3, V, D_2) *_{\mathcal{MCG}_+(S^3,V,D_2,D_2')} \mathcal{MCG}_+(S^3, V, D_2 \cup D_2').$$

Regarding $N(E; E(V))$ as a product $E \times [0,1]$, set $E_+ = E \times \{1\}$ and $E_- = E \times \{0\}$. Set $\alpha = \partial D_2 \setminus (\partial E \times \partial I)$ and $\alpha' = \partial D_2' \setminus (\partial E \times \partial I)$. Recall that $V \cup (E \times [0,1])$ and $N(K_1)$ are ambient isotopic.

Let p be a dot of $(E(K_1), \partial N(K_1))$. By Proposition 1.4, we have $\mathcal{MCG}(S^3, N(K_1), p) \cong \mathcal{MCG}(E(K_1), p)$. Let $P : \pi_1(\partial N(K_1), p) \to \mathcal{MCG}(E(K_1), p)$ be a point-pushing map. The map P induces a map $\bar{P} : \pi_1(\partial N(K_1), p)/\ker P \to \mathcal{MCG}(E(K_1), p)$. Then there exists the Birman exact sequence

$$1 \to \pi_1(\partial N(K_1), p)/\ker P \xrightarrow{\bar{P}} \mathcal{MCG}(E(K_1), p) \to \mathcal{MCG}(E(K_1)) \to 1.$$

Since $\pi_1(\partial N(K_1), p)$ is an abelian group, it is obvious that the group $\bar{P} : \pi_1(\partial N(K_1), p)/\ker P$ is finitely presented. Hence $\mathcal{MCG}(E(K_1), p)$ is finitely generated by Lemma 1.1.

$$\begin{aligned}\mathcal{MCG}_+(S^3, V, D_2) &\cong \mathcal{MCG}_+(S^3, V \cup (E \times [0,1]), E_+ \cup \alpha \cup E_-)\\ &\cong \mathcal{MCG}_+(S^3, N(K_1), p) \rtimes \mathbb{Z}\\ &\cong \mathcal{MCG}_+(E(K_1), p) \rtimes \mathbb{Z},\end{aligned}$$

i.e. $\mathcal{MCG}_+(S^3, V, D_2)$ is an extension of $\mathcal{MCG}_+(E(K_1), p)$ by $\mathbb{Z}$. Therefore the group $\mathcal{MCG}_+(S^3, V, D_2)$ is finitely presented due to Lemma 1.1.

Let D be a meridian disk of $N(K_1)$. Set $A = N(\partial D; \partial N(K_1))$. Assume that the dot p is contained in $\mathrm{Int}\, A$.

$$\begin{aligned}\mathcal{MCG}_+(S^3, V, D_2 \cup D_2') &\cong \mathcal{MCG}_+(S^3, V \cup (E \times [0,1]), E_+ \cup \alpha \cup E_- \cup \alpha')\\ &\cong \mathcal{MCG}_+(S^3, N(K_1), A, p) \rtimes (\mathbb{Z}/2\mathbb{Z}),\\ \mathcal{MCG}_+(S^3, V, D_2, D_2') &\cong \mathcal{MCG}_+(S^3, V \cup (E \times [0,1]), E_+ \cup \alpha \cup E_- \cup \alpha', \alpha)\\ &\cong \mathcal{MCG}_+(S^3, N(K_1), A, p).\end{aligned}$$

Hence $\mathcal{MCG}_+(S^3, V, D_2 \cup D_2')$ and $\mathcal{MCG}_+(S^3, V, D_2, D_2')$ are finitely generated as well by Lemmas 1.1 and 2.6.

Thus we obtain the following theorem.

Theorem 3.12. *If K_1 is non-trivial and K_2 is trivial, then the symmetry group $\mathcal{MCG}(S^3, V)$ of V is finitely presented.*

Example. Consider the case where K_1 is a trefoil and K_2 is the unknot as was depicted in Figure 6. It is well known that $\mathcal{MCG}_+(S^3, N(K_1))$ is homomorphic to the group $\mathbb{Z}/2\mathbb{Z}$ generated by an automorphism f shown in Figure 7, see e.g. [21].



FIGURE 7. The generator f of $\mathcal{MCG}(S^3, N(K_1))$.



(i) (ii)

(iii) (iv)

FIGURE 8. Automorphisms g_1, g_2, g_3 and g_4.

Let D_2 and D_2' be as in the above argument. We may compute the vertex and edge stabilizers of the tree $\mathcal{P}'(V)$ as follows:

$$\begin{aligned}\mathcal{MCG}_+(S^3, V, D_2) &= \langle g_1, g_2, g_3 \mid {g_1}^2, {g_2}^6, g_1 g_2 {g_1}^{-1} g_2, g_1 g_3 {g_1}^{-1} {g_3}^{-1}, g_2 g_3 {g_2}^{-1} {g_3}^{-1} \rangle \\ &\cong ((\mathbb{Z}/6\mathbb{Z}) \rtimes (\mathbb{Z}/2\mathbb{Z})) \ltimes \mathbb{Z},\end{aligned}$$

$$\begin{aligned}\mathcal{MCG}_+(S^3, V, D_2 \cup D_2') &= \langle g_1, g_2, g_4 \mid {g_1}^2, {g_2}^6, {g_4}^2 {g_2}^{-1}, g_1 g_4 {g_1}^{-1} g_4 \rangle \\ &\cong (\mathbb{Z}/12\mathbb{Z}) \rtimes (\mathbb{Z}/2\mathbb{Z}),\end{aligned}$$

$$\mathcal{MCG}_+(S^3, V, D_2, D_2') = \langle g_1, g_2 \mid {g_1}^2, {g_2}^6,\ g_1 g_2 {g_1}^{-1} g_2 \rangle \cong (\mathbb{Z}/6\mathbb{Z}) \rtimes (\mathbb{Z}/2\mathbb{Z}).$$

Hence, we have

$$\begin{aligned}\mathcal{MCG}_+(S^3,V) &= \mathcal{MCG}_+(S^3,V,D_2) *_{\mathcal{MCG}_+(S^3,V,D_2,D_2')} \mathcal{MCG}_+(S^3,V,D_2\cup D_2') \\ &= \langle g_1,g_3,g_4 \mid {g_1}^2, {g_4}^{12}, g_1g_3{g_1}^{-1}{g_3}^{-1}, g_1g_4{g_1}^{-1}g_4, g_3g_4{g_3}^{-1}{g_4}^{-1}\rangle.\end{aligned}$$

Combining Theorems 1.6, 3.12 and Proposition 3.3 we deduce the following conclusive theorem for the case of reducible handlebody-knots of genus two.

Theorem 3.13. *Let $V \subset S^3$ be a reducible handlebody-knot of genus two. Then the symmetry group $MCG(S^3,V)$ is finitely presented.*

4. The symmetry groups of irreducible handlebody-knots of genus two

A handlebody-knot of genus at least two is said to be *hyperbolic* if its exterior admits a complete hyperbolic structure with geodesic boundary.

Due to Thurston's hyperbolization theorem [20] and an equivariant torus theorem [17] it follows that a compact, orientable, irreducible, ∂-irreducible 3-manifold admits a hyperbolic structure with geodesic boundary.

As was remarked in Section 1.1, if V is an irreducible handlebody-knot of genus two, then $E(V)$ is ∂-irreducible. Hence we have the following lemma.

Lemma 4.1. *An irreducible handlebody knot V of genus two is hyperbolic if and only if $E(V)$ contains neither essential tori nor essential annuli.*

Proposition 4.2. *If a handlebody-knot V is hyperbolic, then $\mathcal{MCG}(S^3,V)$ is finite.*

Proof. The mapping class group $\mathcal{MCG}_+(E(V))$ is finite due to the well-known fact that the isometry group of a compact hyperbolic manifold is finite, see [22]. It follows from Lemma 1.2 that $\mathcal{MCG}_+(S^3,V)$ is also finite, hence so $\mathcal{MCG}(S^3,V)$ is. □

Let $V \subset S^3$ be a handlebody-knot of genus two. In the following, we always assume that V is irreducible. By Lemma 4.1 and Proposition 4.2, it remains to consider the case where $E(V)$ contains an essential torus or an essential annulus.

Let $V_0 \subset S^3$ be a trivial handlebody-knot of genus two. A non-separating simple closed curve is said to be *primitive* if it bounds a primitive disk of V_0. A separating simple closed curve is said to be *primitive* if it bounds disks in both V_0 and $E(V_0)$.

Lemma 4.3. *Let $V_0 \subset S^3$ be a trivial handlebody-knot of genus two. Let A_0 be a non-separating annulus properly embedded in V_0. Set $\partial A_0 = C_0 \cup C_1$. Suppose that A_0 is compressible and that C_0 (say) is not primitive. Then $\mathcal{MCG}(S^3,V_0,A_0)$ is a finite group.*

Proof. We compress the annulus A_0 along a compression disk and denote the resulting disks by D_0 and D_1, where $\partial D_0 = C_0$ and $\partial D_1 = C_1$. Then it follows that $\mathcal{MCG}(S^3,V_0,A_0)$ is a subgroup of $\mathcal{MCG}(S^3,V_0,D_0,D_1)$. Since D_0 is not primitive, $\mathcal{MCG}(S^3,V_0,D_0,D_1)$ is a finite group due to [7, Proposition 17.2]. □

An annulus A properly embedded in $E(V)$ is called an *unknotting annulus* of V if $V \cup N(A;E(V))$ is a trivial handlebody-knot of genus two.

Theorem 4.4. *Let A be an unknotting annulus of V and set $V_0 = V \cup N(A; E(V))$. Suppose that A is a unique unknotting annulus of V up to ambient isotopy. Let C be the core $S^1 \times \{1/2\}$ of A. Regard $N(A; E(V))$ as $A \times [0, 1]$ and set $A_0 = C \times [0, 1] \subset N(A; E(V))$. If A_0 is compressible in V_0, then $\mathcal{MCG}(S^3, V)$ is a finite group.*

Proof. It is clear from the definition that A_0 is non-separating in V_0. Set $\partial A_0 = C_0 \cup C_1$. Since A_0 is compressible, both C_0 and C_1 bound disks D_0 and D_1 in V_0, respectively. Then there exists a simple arc l embedded in V_0 such that

(1) $l \cap (D_0 \cup D_1) = \partial l$;
(2) $l \cap D_0 \neq \emptyset$ and $l \cap D_1 \neq \emptyset$; and
(3) $\overline{N(l \cup D_0 \cup D_1; V_0) \setminus \partial V_0}$ consists of three components D'_0, D'_1 and A'_0, where D'_0 is parallel to D_0, D'_1 is parallel to D_1 and A'_0 is parallel to A_0.

See Figure 9. Since A_0 is non-separating, either C_0 or C_1 is non-separating. Assume that both C_0 and C_1 are primitive non-separating simple closed curves. Recall that the handlebody-knot V is obtained by cutting V_0 along an annulus A'_0. Then we may isotopy V_0 in S^3 as shown in Figure 10. The resulting figure implies that V reducible, which contradicts our assumption. (In fact, V is obtained from an unknotted solid torus by digging a tunnel, it follows from [13] that V is an unknotted handlebody of genus two.) We may also deduce analogously a contradiction in the case where C_0 is a primitive separating simple closed curve and C_1 is a primitive non-separating one. As a consequence, A_0 satisfies the assumption of Lemma 4.3.



FIGURE 9. Primitive disks D_0 and D_1 and an arc l connecting them.



FIGURE 10. Cutting V_0 along A'_0 and an ambient isotopy.

Let f be an element of $\mathcal{MCG}(S^3, V)$. Since A is a unique unknotting annulus for V, we may assume that $f(N(A; E(V))) = N(A; E(V))$. Hence f uniquely determines an element $\tilde{f}$ of $\mathcal{MCG}(S^3, V_0, A)$. Conversely, each element of $\mathcal{MCG}(S^3, V_0, A)$ uniquely determines an element of $\mathcal{MCG}(S^3, V)$. Thus we have $\mathcal{MCG}(S^3, V) \cong \mathcal{MCG}(S^3, V_0, A_0)$ and the assertion follows immediately from Lemma 4.3. □

The Svarć-Milnor-Lemma and the proof of Theorem 4.4 allow us to deduce the following corollary:

Corollary 4.5. *Suppose that* V *admits finitely many unknotting annuli* $A_1, A_2, \dots, A_n$ *up to ambient isotopy. Let* C^i *be the core of* A_i*. Regard* $N(A_i; E(V))$ *as* $A_i \times [0,1]$ *and set* $A_i^d = C^i \times [0,1] \subset N(A_i; E(V))$*. If* A_i^d *is compressible in* $V \cup N(A_i; E(V))$ *for each* $i = 1, 2, \dots, n$*, then* $\mathcal{MCG}(S^3, V)$ *is a finite group.*

Example. The left-hand side of Figure 11 illustrates a series of handlebody-knots V_n ($n \geqslant 3$) of genus two. We note that in the table of irreducible handlebody-knots in [19], $V_3 = 5_1$ and $V_4 = 6_1$. It follows easily from the argument in [24, 25] that these handlebody-knots are irreducible and that the annulus A depicted in the figure is a unique unknotting annulus properly embedded in $E(V_n)$. Moreover, it follows that all of these handlebody-knots are not *amphicheiral*, i.e. $\mathcal{MCG}(S^3, V_n) \cong \mathcal{MCG}_+(S^3, V_n)$. The right-hand side of the figure shows the curves C_0 and C_1 in the case $n = 3$. The "hyperelliptic" involution is the unique non-trivial element of $\mathcal{MCG}(S^3, V_0)$ which preserve the curves $C_0 \cup C_1$. However, this involution does not preserve the annulus A_0, hence we have $\mathcal{MCG}(S^3, V_n) = 1$.



FIGURE 11. The irreducible handlebody-knot V_n and the curves C_0 and C_1.

Acknowledgements. The author wishes to express his gratitude to Sangbum Cho for his helpful comments on the preliminary version of the paper. Part of this work was carried out while the author was visiting the Mathematisches Forschungsinstitut Oberwolfach. The institute kindly offered an invitation for the stay while his University was affected by the 2011 Tohoku earthquake. He is grateful to the MFO and its staffs for the offer, the financial support and warm hospitality. He also would like to thank the Leibniz Association for their travel support. This work is supported by Grant-in-Aid for Young Scientists (B) 20525167.

REFERENCES

[1] Akbas, E., A presentation for the automorphisms of the 3-sphere that preserve a genus two Heegaard splitting, Pacific J. Math. **236** (2008), 201–222.
[2] Benedetti, R., Frigerio, R., Levels of knotting of spatial handlebodies, arXiv:1101.2151.
[3] Birman, J. S., *Braids, links and mapping class groups*, Ann. Math. Stud. **82**, Princeton Univ. Press, Princeton, N.J., 1974.

[4] Brown, K., Presentations for groups acting on simply-connected complexes, J. Pure Appl. Algebra **32** (1984) 1–10.

[5] Cho, S., Homeomorphisms of the 3-sphere that preserve a Heegaard splitting of genus two, Proc. Amer. Math. Soc. **136** (2008), 1113–1123.

[6] Cho, S., Goeritz groups for lens spaces, *preprint.*

[7] Cho, S., McCullough, D., The tree of knot tunnels, Geom. Topol. **13** (2009), 769–815.

[8] Cooper, D., Hodgson C., Kerkhoff, S., Three-dimensional orbifolds and cone-manifolds. MSJ Mem. **5**, 2000.

[9] Flapan, E., Naimi, R., Pommersheim, J., Tamvakis, H., Topological symmetry groups of embedded graphs in the 3-sphere, Comment. Math. Helv. **80** (2005), 317–354.

[10] Johannson, K., *Homotopy equivalences of* 3*-manifolds with boundaries*, Lecture Notes in Math., **761**, Springer, Berlin, 1979.

[11] Fomenko, A. T., Matveev, S. V., *Algorithmic and Computer Methods for Three-Manifolds*, Math. Appl. **425**, Kluwer Acad. Publ., Dordrecht, 1997.

[12] Goeritz, L., Die Abbildungen der Brezelfläache und der Vollbrezel vom Geschlecht 2. Abh. Math. Semin. Hamb. Univ. **9** (1933), 244–259.

[13] Gordon, C., On primitive sets of loops in the boundary of a handlebody. Topol. Appl. **27** (1987), 285–299.

[14] Gordon, C., Luecke, J., Knots Are Determined by Their Complements, J. Amer. Math. Soc. **2** (1989), 371–415.

[15] Hamenstädt, U., Hensel, S., The geometry of the handlebody groups I: Distortion, arXiv:1101.1838v2.

[16] Heard, D., Hodgson, C., Martelli, B., Petronio, C., Hyperbolic Graphs of Small Complexity, Trans. Amer. Math. Soc. **326** (1991), 887–906. Experiment. Math. **19** (2010), 211–236.

[17] Holzmann, W. H., An equivariant torus theorem for involutions, Trans. Amer. Math. Soc. **326** (1991), 887–906.

[18] Ishii, A., Moves and invariants for knotted handlebodies, Algebr. Geom. Topol. **8** (2008), 1403–1418.

[19] Ishii, A., Kishimoto, K., Moriuchi, H., Suzuki, M., A table of genus two handlebody-knots up to six crossings, *preprint.*

[20] Kapovich, *Hyperbolic manifolds and discrete groups*, Modern Birkhauser Classics, Birkhäuser Boston, Boston, 2001.

[21] Kawauchi, A., *Survey of Knot Theory*, Birkhäuser, Basel, 1996.

[22] Kojima, S., Isometry transformations of hyperbolic 3-manifolds, Topology Appl. **29** (1988), 297–307

[23] McCullough, D., Virtually geometrically finite mapping class groups of 3-manifolds, J. Diff. Geom. **33** (1991), 1–65.

[24] Motto, M., Inequivalent genus 2 handlebodies in S^3 with homeomorphic complement, Topology Appli. **36** (1990), 283–290.

[25] Lee, J. H., Lee, S., Inequivalent handlebody-knots with homeomorphic complements, *preprint.*

[26] Scharlemann, M., Automorphisms of the 3-sphere that preserve a genus two Heegaard splitting, Bol. Soc. Mat. Mexicana **10** (2004), pp503–514.

[27] Serre, J.-P., *Arbres, amalgames,* SL_2, Astérisque **46**, Société Mathématique de France, Paris, 1977.

[28] Simon, J, Topological chirality of certain molecules, Topology **25** (1986), 229–235.

[29] Sims, C. C., *Computation with Finitely Presented Groups*, Encyclopedia of Mathematics and its Applications, Cambridge Univ. Press, Cambridge, 1994.

Mathematical Institute, Tohoku University, Sendai, 980-8578, JAPAN
E-mail address: `koda@math.tohoku.ac.jp`